



*The Alberta Lake Management Society
Volunteer Lake Monitoring Program*

Cooking Lake

• • • • • • • • • •

2007 Report

Completed with support from:





*The Alberta Lake Management Society
Volunteer Lake Monitoring Program*

• • • • • • • • • •

Alberta Lake Management Society

Address: PO Box 4283, Edmonton,
Alberta, T6E 4T3

Phone: 780-702-ALMS

E-mail: info@alms.ca

Water is integral to supporting and maintaining life on this planet as it moderates the climate, creates growth and shapes the living substance of all of Earth's creatures. It is the tide of life itself, the sacred source. David Suzuki (1997). The Sacred Balance.

Alberta Lake Management Society's Lakewatch Program

Lakewatch has several important objectives, one of which is to collect and interpret water quality on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. Lakewatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in Lakewatch and readers requiring more information are encouraged to seek these sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the Lakewatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.

Acknowledgements

The Lakewatch program is made possible through the dedication volunteers and the Lakewatch Chairs, Théo Charette and Ron Zurawell. Gary Stuart and his granddaughter, Olivia contributed their time and collected field data during 2007. Numerous Alberta Environment staff also contributed to successful completion of the 2007 program. We would like to thank Jill Anderson and Wendy Markowski who were summer interns with ALMS in 2007. Project Technical Coordinator, Megan McLean was instrumental in planning and organizing the field program. Technologists, Mike Bilyk, Brian Jackson and John Willis were involved in the logistics planning and training aspects of the program. Doreen LeClair was responsible for data management. Théo Charette, Ron Zurawell (Limnologist, AENV), and Lori Nuefeld prepared the original report, which was updated by Heather Powell for the 2007 report. The Lakewatch program was financially supported by Alberta Environment and the Lakeland Industry and Community Association (LICA).

Cooking Lake

Introduction

Cooking Lake is a large, shallow lake located in the North Saskatchewan River Basin, about 25 km east of the city of Edmonton. The nearest towns are South Cooking Lake, North Cooking Lake and Collingwood Cove.



Figure 1. Photo of Cooking Lake taken from the Atlas of Alberta Lakes.

The lake's name is a translation of the Cree *opi-mi-now-wa-sioo*, which means "a cooking place". It is thought that the shores of Cooking Lake were a favorite Cree campground (Mitchell and Prepas 1990). South Cooking Lake is the oldest settlement on the lake while North Cooking Lake became an important recreation area after 1909, when the Grand Trunk Pacific Railway line from Edmonton was completed. Special weekend trains from Edmonton brought tourists to the north end of the lake, and passenger boats transported the visitors to sandy beaches on the south shore (Mitchell and Prepas 1990, **Figure 1**).




Figure 2. Bathymetric map of cooking lake. Each line represents a 1.5 m change in water depth.

Cooking lake is a relatively large lake with a surface area of 36 km². Despite the lake's large size it is very shallow with a mean depth of 1.7 m and a maximum depth of 4.6 m the last time soundings were taken (**Figure 2**). The water in Cooking Lake is very nutrient rich, and dense algal blooms occur from mid- to late summer. Consequently, there are few recreational facilities at the lake. The lake is most popular for wind surfing, sailing, power boating and bird watching. It is considered an important breeding and migration stopover area for waterfowl. No sport fish live in the lake because they cannot survive the low oxygen conditions that occur during the winter (Mitchell and

Prepas 1990).

Results

Water Levels

Beginning in the early 1970s water levels in Cooking Lake have been monitored by Environment Canada under the joint federal-provincial hydrometric agreement. Water levels are measured as the elevation in meters above sea level (m asl) of the surface of the lake. Between the mid 1970's and mid 1990s the water level of Cooking Lake has declined by roughly 0.5 m (**Figure 3**). This decline in water level is particularly concerning because Cooking Lake was shallow to begin with. Measurements of recent water levels were unavailable for Cooking Lake however Lakewatch volunteers reported that the maximum depth of the lake during sampling in the summer of 2006 was 1.03 m, which suggests a further decline in water levels.




Figure 3. Water levels for Cooking Lake measured as meters above sea level (m asl) of the surface of the lake.

Measurements of recent water levels were unavailable for Cooking Lake however Lakewatch volunteers reported that the maximum depth of the lake during sampling in the summer of 2006 was 1.03 m, which suggests a further decline in water levels.

Water Temperature and Dissolved Oxygen

Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. Please refer to the end of this report for descriptions of technical terms.

Cooking Lake is a shallow, polymictic lake, which means that the water column mixes many times throughout the summer. Because of frequent mixing the water temperature and dissolved oxygen concentration is relatively the same at all water depths of the lake (**Figure 4**).

On 27 July and 9 August, dissolved oxygen (DO) concentrations in Cooking Lake were < 5 mg/L (**Figure 4**). At all other dates, oxygen levels in surface layers of Cooking Lake were within the acceptable range for surface water quality, according to Alberta Environment guidelines (DO \geq 5.0 mg/L). Wind-induced mixing of the lake does not occur once winter ice covers the lake, and oxygen concentrations in the lake decline (data not shown). The low oxygen concentration in the winter makes it difficult for sport fish to survive in Cooking Lake.




Figure 4. Water temperature (°C) and dissolved oxygen (mg /L) profiles for Cooking Lake during the summer and autumn of 2007.

Water clarity and Secchi Depth

Water clarity is influenced by suspended materials, both living and dead, as well as dissolved colored compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring but then becomes more turbid with increased algal growth as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi disk depth.

Cooking Lake is a shallow, polymictic lake and as such is often turbid (e.g. murky). Although light penetrated to ~50% of the total lake depth (average Secchi disk depth = 0.51 m) during the summer of 2007, the lake appeared turbid. Because the lake is shallow and light penetrates through most of the water column, algae can grow throughout the lake. Thus, as algae become more abundant, water clarity decreases. This is evidenced by maximum water clarity (Secchi disk depth = 0.6 m on 27 July) when algal biomass was low. Minimum water clarity (Secchi disk depth = 0.45 m on 26 August) corresponded to a mid-summer peak in algal biomass (Chla = 59.3 mg/m³).

Water chemistry

Based on lake water characteristics, Cooking Lake is considered to be hypereutrophic (very nutrient rich; see *A Brief Introductory to Limnology* at the end of this report) (**Table 1**). Lakes in Alberta are naturally productive. However Cooking Lake is high in nutrient concentrations and algae biomass compared to other Alberta lakes. Phosphorous concentration peaked in August and nitrogen increased between July and October

(**Figure 5**). The amount of algae in the lake (chlorophyll *a*) peaked in early August in response to increased nutrient levels (**Figure 5**).




Figure 5. Total phosphorus, total nitrogen, and chlorophyll *a* (a measure of algae biomass) concentrations for Cooking Lake during the summer and autumn of 2007.

Cooking Lake is well-protected from acidification; its pH of 8.9 is well above that of pure water (i.e., pH 7). Cooking Lake is a hardwater, slightly saline lake. Ion concentrations are relatively high (**Table 1**). Dominant ions include bicarbonate, carbonate, sodium, and magnesium. Ions are supplied by weathering in the watershed, groundwater inflows, and from deposition of pollutants from the air. Reduced water levels can also cause an increase in the concentration of ions as water evaporates, leaving the ions behind to accumulate. Declining water levels, due to a 20-year drought in the region, may be responsible for some of the high ion concentrations.

The average concentrations of various heavy metals (as total recoverable concentrations) in Cooking Lake were not available for the summer of 2007, with the exception of iron. Iron concentrations and CCME heavy metal guidelines for the Protection of Freshwater Aquatic Life are presented for reference in **Appendix 1**.

Table 1. Mean water chemistry values for Cooking Lake, Alberta summer 2007.

Parameter	2007
TP ($\mu\text{g/L}$)	402.6
Total dissolved P ($\mu\text{g/L}$)	238.4
Chlorophyll- <i>a</i> ($\mu\text{g/L}$)	49.6
Secchi disk depth (m)	0.51
TN (mg/L)	9.3
NO ₂₊₃ ($\mu\text{g/L}$)	20.7
NH ₄ ($\mu\text{g/L}$)	185.2
Dissolved organic C (mg/L)	98.2
Ca (mg/L)	21.0
Mg (mg/L)	76.1
Na (mg/L)	542
K (mg/L)	73.2
SO ₄ (mg/L)	744.3
Cl (mg/L)	54.5
CO ₃ (mg/L)	63.7
HCO ₃ (mg/L)	730
pH	8.9
Conductivity ($\mu\text{S/cm}$)	2730
Total dissolved solids (mg/L)	1933
Total Alkalinity (mg/L CaCO ₃)	705

Note: TP = total phosphorus, Chl_a = chlorophyll *a*, TN = total Kjeldahl nitrogen, NO₂₊₃ = nitrate+nitrite, NH₄ = ammonium, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO₄ = sulphate, Cl = chloride, CO₃ = carbonate, HCO₃ = bicarbonate.
 *Atlas of Alberta Lakes (Mitchell and Prepas, 1990).

References

- Mitchell, P. and E. Prepas. 1990. Atlas of Alberta Lakes. University of Alberta Press.
- Vollenweider, R.A., and J. Kerekes, J. 1982. Eutrophication of Waters. Monitoring, Assessment and Control. Organization for Economic Co-Operation and Development (OECD), Paris. 156p.
- Welch, E.B. 1980. Ecological Effects of Waste Water. Cambridge University Press.

Appendix 1

Mean concentrations of metals in Cooking Lake were unavailable for the summer of 2007, with the exception of iron. The CCME heavy metal Guidelines for the Protection of Freshwater Aquatic Life (unless otherwise indicated) are presented for reference.

Metals (total)	2007	Guidelines
ALUMINUM µg/L	-	100 ^a
ANTIMONY µg/L	-	6 ^e
ARSENIC µg/L	-	5
BARIUM µg/L	-	1000 ^e
BERYLLIUM µg/L	-	100 ^{d,f}
BISMUTH µg/L	-	-
BORON µg/L	-	5000 ^{e,f}
CADMIUM µg/L	-	0.085 ^b
CHROMIUM µg/L	-	-
COBALT µg/L	-	1000 ^f
COPPER µg/L	-	4 ^c
IRON µg/L	88.6	300
LEAD µg/L	-	7 ^c
LITHIUM µg/L	-	2500 ^g
MANGANESE µg/L	-	200 ^g
MOLYBDENUM µg/L	-	73 ^d
NICKEL µg/L	-	150 ^c
SELENIUM µg/L	-	1
SILVER µg/L	-	0.1
STRONTIUM µg/L	-	-
THALLIUM µg/L	-	0.8
THORIUM µg/L	-	-
TIN µg/L	-	-
TITANIUM µg/L	-	-
URANIUM µg/L	-	100 ^e
VANADIUM µg/L	-	100 ^{f,g}
ZINC µg/L	-	30
FLUORIDE mg/L	-	1.5

With the exception of fluoride (which reflects the mean concentration of dissolved fluoride only), values represent means of total recoverable metal concentrations.

^a Based on pH ≥ 6.5; calcium ion concentration [Ca⁺²] ≥ 4 mg/L; and dissolved organic carbon concentration [DOC] ≥ 2 mg/L.

^b Based on water Hardness of 300 mg/L (as CaCO₃).

^c Based on water Hardness > 180 mg/L (as CaCO₃).

^d CCME interim value.

^e Based of Canadian Drinking Water Quality guideline values.

^f Based of CCME Guidelines for Agricultural Use (Livestock Watering).

^g Based of CCME Guidelines for Agricultural Use (Irrigation).

A Brief Introduction to Limnology

Indicators of water quality

The goal of **Lakewatch** is to collect water samples necessary to determine the water quality of lakes. Though not all encompassing, the variables measured in **Lakewatch** are sensitive to human activities in watersheds that may cause impacts to water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are affected (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded fish habitat and production of noxious odors. Large increases in nutrients over time may also indicate sewage inputs, which in turn, may result in other human health concerns such as harmful bacteria or protozoans (e.g. *Cryptosporidium*).

Temperature and mixing

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality (Figure 5). Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. A third layer, known as the metalimnion, provides an effective barrier between the epi- and hypolimnion. The metalimnion reflects a rapid transition in water temperature known as the **thermocline**. A thermocline typically occurs when water temperature changes by several degrees within one-meter of depth. The thermocline acts as an effective physico-chemical barrier to mixing between the hypolimnion and epilimnion, restricts downward movement of elements, such as oxygen, from the surface into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point the water mixes from top to bottom in what is called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and 0° C water on the top.




Figure 5: Difference in the circulation of the water column depending on thermal stratification.

In spring another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are termed **polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic** meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice-free season the lake is polymictic.

Dissolved Oxygen

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the hypolimnion. At the same time oxygen is depleted in the hypolimnion by decomposition. The result is that the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill, which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines state dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg/L and should not average less than 6.5 mg/L over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg/L in areas where early life stages of aquatic biota, particularly fish, are present.

General Water Chemistry

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called ions. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. Hydrophobic (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

Phosphorus and Nitrogen

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits terrestrial plants and plants and algae of tropical lakes, phosphorus is usually in shortest supply in temperate lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When pollution originating from livestock manure and human sewage enters lakes not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

Chlorophyll-a

Chlorophyll-a is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll-a can be easily extracted from algae in the laboratory. Consequently, chlorophyll-a is a good estimate of the amount of algae in the water. Larger aquatic plants, known as macrophytes, rather than algae, dominate some highly productive lakes. In these lakes, chlorophyll-a and nutrient values taken from water samples do not include productivity from large aquatic plants. As a result, lakes like Chestermere, which are dominated by macrophytes, reflect lower-nutrient trophic states than would otherwise result if macrophyte-based chlorophyll were included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

Secchi Disk Depth

Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. Secchi disk depth is the oldest, simplest, and quickest quantitative measure of water clarity. A Secchi disk is a black and white disk that is lowered down through the water column until it can no longer be seen. Secchi disk depth is the midpoint between the depth at which it disappears when lowered and reappears when it is pulled up again. The Secchi disk depth in lakes with high algal biomass will generally be low. Secchi disk depth, however, is not only affected by algae, high concentrations of suspended sediments, particularly fine clays or glacial till common in plains or mountain reservoirs of Alberta, also impact water clarity. Mountain reservoirs may have exceedingly shallow Secchi disk depths despite low algal growth and nutrient concentrations.

The euphotic zone, calculated as twice the Secchi disk depth, is the portion of the water column that has sufficient light for aquatic plants to grow. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Aquatic plants are important because they ensure clear lake water by reducing shoreline erosion and stabilizing lake bottom sediments. Many lakes in Alberta are shallow and have bottom sediments with high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment-laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and fish, depend on aquatic plants for food and shelter.

Trophic State

Trophic state is a classification system for lakes that depends on fertility and is a useful index for rating and comparing lakes. From low to high nutrient and algal biomass (as chlorophyll-a) concentrations, the trophic states are: oligotrophic, mesotrophic, eutrophic and hypereutrophic. The nutrient and algal biomass concentrations that define these categories are shown in Table 2

and a graph of Alberta lakes compared by trophic state can be found on the ALMS website. A majority of lakes in Alberta are meso- to eutrophic because they naturally contain high nutrient concentrations due to our deep fertile soils. Thus, lakes in Alberta are susceptible to human impacts because they are already nutrient-rich; any further nutrient increases can bring about undesirable conditions illustrated in Figure. 6.




Figure 6: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980.

Table 2: Trophic status based on lake water characteristics

Trophic state	Total Phosphorus (µg/L)	Total Nitrogen (µg/L)	Chlorophyll a (µg/L)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 - 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 - 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1

Note: These values are from a detailed study of global lakes reported in Nurnberg, 1996. Alberta Environment uses slightly different values for TP and CHL based on those of the OECD reported by Vollenweider and Kerekes (1982). The AENV and OECD cutoffs for TP are 10, 35 and 100; for CHL are 3, 8 and 25. AENV does not have TN or Secchi depth criteria. The corresponding OECD exists for Secchi depth and the cutoffs are 6, 3 and 1.5 m.